

Remarks

In view of the above amendments and the following remarks, reconsideration of the outstanding office action is respectfully requested.

Claims 1, 3, 9, and 11 have been amended, claim 2 has been cancelled without prejudice, and new claim 53 has been added. Descriptive support for the claim 1 amendments is found, *inter alia*, in original claim 2, paragraphs [0038], [0039], [0115] with Table 1, [0130], [0132], [0137], [0139] and [0144]. Descriptive support for claim 9 amendments can be found in paragraphs [0042], [0054], and Example 1. Amendments to claim 11 are supported by paragraphs [0043] and [0045]. Descriptive support for new claim 53 is provided in claim 1 and paragraph [0039]. No new matter has been added by way of the above amendments.

Claims 1, 3-6, 9, 11-13, 19-22 and 53 are under examination, and claims 14-18, 23-26, 32-46, and 48-52 stand withdrawn. No excess claim fees are due with this submission.

The objection to claim 11 for informalities is overcome by the above amendments. The objection to claim 11 should be withdrawn.

The rejection of claims 1, 2, 6, 9, 11-13, and 19-22 under 35 U.S.C. § 112, first paragraph, for failing to comply with the written description requirement is respectfully traversed.

The United States Patent and Trademark Office ("PTO") has asserted that the application fails to identify the structure/function relationship of factor VIII proteins and therefore cannot support the genus of factor VIII proteins as claimed. Applicants respectfully disagree, because the specification demonstrates that the structure/function of factor VIII proteins was well known in the art, and factor VIII nucleic acid and amino acid sequences were well known in the art at the time of filing.

As evidence that the structure and function of factor VIII proteins was well known in the art, the specification identifies wild-type factor VIII nucleic acid and amino acid sequences from human, rat, mouse, dog, chimp, and pig by their Genbank accessions. These Genbank accessions are incorporated by reference into the specification. The specification also references the availability of homology and alignment analyses for factor VIII proteins via the HAMSTeRS internet site (see paragraph [0033]). Attached as Exhibit 1 is the referenced

HAMSTeRS alignment of human, porcine, murine, and canine factor VIII, which demonstrates the high degree of homology among these proteins that was known in the art at the time of filing.

With respect to the human sequence, which for the above-noted reasons represents the genus of mammalian factor VIII, the sequences and domains are explicitly identified in paragraphs [0034] – [0038]. The presence of a conserved A1 domain calcium binding site is identified in paragraph [0038] and illustrated in Figure 6, which shows an alignment of corresponding regions of human, murine, porcine, and canine factor VIII.

Moreover, the specification explicitly identifies mutant factor VIII proteins that were previously known in the art and can be modified in accordance with the present invention. These are identified in paragraphs [0041] – [0054], and the contents of cited references are incorporated by reference into the application.

From the foregoing, persons of skill in the art fully appreciated at the time of filing that a known structure/function relationship exists for mammalian factor VIII proteins, whether a wild-type form or a previously known mutant form.

University of California v. Eli Lilly and Co., 119 F.3d 1559, 43 USPQ2d 1398 (Fed. Cir. 1997) is cited by the PTO for the proposition that a general recitation of function is insufficient to define the structure of a protein. This decision confirmed that the single disclosed rat cDNA encoding insulin did not support claims to human cDNA encoding insulin (no species disclosed), mammalian cDNA encoding insulin (1 species disclosed) and vertebrate cDNA encoding insulin (1 species disclosed). However, in sharp contrast to the facts in *University of California*, as noted above many mammalian factor VIII proteins were known in the art prior to the present invention — both wild-type and mutant forms. Importantly, their structural homology was also known (see attached Exhibit 1 and Figure 6 of application). Applicants therefore submit that the recitation and incorporation by reference of six different wild-type mammalian factor VIII proteins and dozens of mutant mammalian factor VIII proteins adequately supports the genus as claimed. The assertion on page 4 of the office action that the specification only supports human factor VIII is therefore improper.

The rejection of claims 9 and 11 is obviated by the above amendments.

For all these reasons, the rejection of claims 1, 2, 6, 9, 11-13, and 19-22 for lack of written description is improper and should be withdrawn.

The rejection of claims 1-6, 9, 11-13, and 19-22 under 35 U.S.C. § 112, second paragraph for indefiniteness is respectfully traversed.

Claims 1 and 6 are rejected for lack of clarity concerning the term “specific activity”. Claim 1 has been amended to recite “specific activity, as measured in a one-stage clotting assay”. This clotting assay is well known in the art and identified in the specification (see Example 10). The one-stage clotting assay measures the ability of a product—in this case, the claimed product—to shorten the clotting time of hemophilia A plasma in a system containing activated partial thromboplastin. It is called a one-stage assay because, after a preincubation of the activated partial thromboplastin, factor VIII-deficient plasma, and the product being tested to achieve contact activation of plasma, the single step of recalcifying the incubation mixture leads to fibrin clot formation. The one-stage clotting assay is described, and compared to other known *in vitro* assays, in Lundblad et al., “Issues with the Assay of Factor VIII Activity in Plasma and Factor VIII Concentrates,” *Thromb. Haemost.* 84:942-948 (2000) (copy attached as Exhibit 2).

Claims 9 and 11 are rejected for lack of clarity in failing to define the term “portion” as it relates to domains A1, A2, A3, C1, and C2. Claim 9 has been amended to instead recite a recombinant factor that is B domainless, which overcomes this basis of rejection. For these reasons, the rejection of claims 1-6, 9, 11-13, and 19-22 for indefiniteness should be withdrawn.

The rejection of claims 1-6, 9, 11-13, and 19-22 under 35 U.S.C. § 102(a) as being anticipated by Wakabayashi et al. “Residues 110-126 in the Factor VIII Heavy Chain Contain a Ca²⁺ Binding Site Required for Cofactor Activity,” *Blood, ASH Annual Meeting Abstracts*, 102(11): p542a, Abstract 1988 (2003) (“Wakabayashi”) is respectfully traversed.

Wakabayashi is not available as prior art under 35 U.S.C § 102(a). Pursuant to the accompanying Declaration of Philip J. Fay under 37 C.F.R. § 1.132 (“Fay Decl.”), Wakabayashi does not evidence knowledge or use of the present invention by others in this country prior to the invention by applicants. Specifically, non-inventors Jan Freas and Qian Zhou prepared and purified reagents under the direction and control of one of the inventors (Fay Decl. ¶4). Neither Jan Freas nor Qian Zhou contributed to the conception of the claimed invention (*Id.*). For these reasons, the rejection of claims 1-6, 9, 11-13, and 19-22 for anticipation by Wakabayashi is improper and should be withdrawn.

The rejection of claims 1, 2, 6, 9, 12, 13, and 19-22 under 35 U.S.C. § 102(b) for anticipation over U.S. Patent No. 5,422,260 to Kaufman et al. ("Kaufman") is respectfully traversed.

The PTO cites Kaufman for disclosing mutant human factor VIII comprising at least one to three amino acid mutations at certain positions and exhibiting procoagulant activity, methods to produce such variants of factor VIII and pharmaceutical compositions of such variants. This rejection is improper, because the PTO has failed to demonstrate that the mutations specified by Kaufman (at amino acid positions 220, 250, 279, 282, 325, 338, 346, 359, 395, 407, 698, 700, 741, 1664, 1680, and 1719) are in or near at least a calcium binding site of the A1 domain. As recited at paragraph [0031], "in or near" means within about five amino acid residues from a residue that directly interacts with Ca^{2+} or Mn^{2+} ions. Also, the PTO has failed to demonstrate that the recombinant factor VIII of Kaufman has a specific activity, as measured by one-stage clotting activity, that is higher than that of a wild-type factor VIII. Given these deficiencies, the rejection of claims 1, 2, 6, 9, 12, 13, and 19-22 for anticipation by Kaufman is improper and should be withdrawn.

The rejection of claims 1, 2, 6, 9, 12, 13, and 19-22 under 35 U.S.C. § 102(e) for anticipation over U.S. Patent Application Publication No. 2005/0100990 to Saenko et al. ("Saenko") is respectfully traversed.

Saenko is cited for teaching methods to increase the procoagulant activity of factor VIII by substituting one or more amino acids in the A2 domain. Saenko further teaches the methods to produce such mutants and pharmaceutical composition comprising such mutant factor VIII. None of these modified residues is present in the A1 domain, and therefore none of these residues involves a calcium binding site of the A1 domain. This rejection is therefore improper. For these reasons, the rejection of claims 1, 2, 6, 9, 12, 13, and 19-22 for anticipation by Saenko should be withdrawn.

The rejection of claim 11 under 35 U.S.C. § 103(a) for obviousness over Saenko in view of U.S. Patent No. 5,859,204 to Lollar et al. ("Lollar") is respectfully traversed.

The teachings and deficiencies of Saenko are noted above. Lollar is cited for teaching a hybrid factor VIII having human and animal factor VIII amino acid sequences, methods of preparation, and use of such hybrid factor VIII. However, PTO has failed to

- 11 -

demonstrate how Loller overcomes the deficiencies of Saenko. For this reason, the rejection of claim 11 for obviousness over the combination of Saenko and Loller is improper and should be withdrawn.

In view of all of the foregoing, applicant submits that this case is in condition for allowance and such allowance is earnestly solicited.

Respectfully submitted,

Date: February 23, 2009

/Edwin V. Merkel/
Edwin V. Merkel
Registration No. 40,087

NIXON PEABODY LLP
1100 Clinton Square
Rochester, New York 14604
Telephone: (585) 263-1128
Facsimile: (585) 263-1600

Exhibit 1: HAMSTeRS Alignment of Human, Porcine, Murine, and Canine Factor VIII

```

HUMFVIII 1 AARRYYLGAVELSDWYMQSDLLSEHLVCEFRPPVPVPSSEPTNTSVVTKKT
PIGFVIII 1 AARRYYLGAVELSDWYMQSDLLSEHLVDTREFFAAGDAPLQISVLYTKT
MURFVIII 1 AARRYYLGAVELSDWYMQSDLLSEHLVDSREFFRMSVPSISLIMVTKT
CANFVIII 1 AARRYYLGAVELSDWYMQSDLLSEHLVDTSFSESRVPGSPILTSVYTKT

HUMFVIII 50 LFVEFTDGLFENIAKPRPPWMGLLGPTIQAEYDITVVITLKNMASHPVSLH
PIGFVIII 51 VFVEFTDQLFSEVARPPFPWMGLLGPTIQAEYDITVVVITLKNMASHPVSLH
MURFVIII 51 VFVEFTDGLFENIAKPRPPFPWMGLLGPTIQAEYDITVVITLKNMASHPVSLH
CANFVIII 51 VFVEFTDGLFENIAKPRPPFPWMGLLGPTIQAEYDITVVITLKNMASHPVSLH

HUMFVIII 100 AVGVSYWKASGEAYEDDQTSQREKEDDKVFPQSSHTYVWQVLKENGPMAS
PIGFVIII 101 AVGVSYFKSSSEGAAYEDDQTSQREKEDDKVFPQSSHTYVWQVLKENGPMAS
MURFVIII 101 AVGVSYWKASGEAYEDDQTSQREKEDDKVFPQSSHTYVWQVLKENGPMAS
CANFVIII 101 AVGVSYWKASGEAYEDDQTSQREKEDDKVFPQSSHTYVWQVLKENGPMAS

HUMFVIII 150 DPPCLTYSYLSHVDLVKDLNSGLIGALLVCREGSLAKEKTQTLHSEFVLLF
PIGFVIII 151 DPPCLTYSYLSHVDLVKDLNSGLIGALLVCREGSLAKEKTQTLHSEFVLLF
MURFVIII 151 DPPCLTYSYLSHVDLVKDLNSGLIGALLVCREGSLAKEKTQTLHSEFVLLF
CANFVIII 151 DPPCLTYSYLSHVDLVKDLNSGLIGALLVCREGSLAKEKTQTLHSEFVLLF

HUMFVIII 200 AVFDEGKSWHSETNDSLTQASARATPKMHTVNGYVNRSLPGLIOCH
PIGFVIII 201 AVFDEGKSWHSETNDSLTQASARATPKMHTVNGYVNRSLPGLIOCH
MURFVIII 201 AVFDEGKSWHSETNDSLTQASARATPKMHTVNGYVNRSLPGLIOCH
CANFVIII 201 AVFDEGKSWHSETNDSLTQASARATPKMHTVNGYVNRSLPGLIOCH

HUMFVIII 250 RKSYYVHWIGMGTTPVHSIFLEOHTFLVRNHRQASLEISPIETLTAQT
PIGFVIII 251 RKSYYVHWIGMGTTPVHSIFLEOHTFLVRNHRQASLEISPIETLTAQT
MURFVIII 251 RKSYYVHWIGMGTTPVHSIFLEOHTFLVRNHRQASLEISPIETLTAQT
CANFVIII 245 RKSYYVHWIGMGTTPVHSIFLEOHTFLVRNHRQASLEISPIETLTAQT

HUMFVIII 300 LMDLGGQFLLFCHISSHQHDMGMEATYKVVDSCPEEPQLRMKNNEEDDYDD
PIGFVIII 301 LMDLGGQFLLFCHISSHQHDMGMEATYKVVDSCPEEPQLRMKNNEEDDYDD
MURFVIII 301 LMDLGGQFLLFCHISSHQHDMGMEATYKVVDSCPEEPQLRMKNNEEDDYDD
CANFVIII 295 LMDLGGQFLLFCHISSHQHDMGMEATYKVVDSCPEEPQLRMKNNEEDDYDD

HUMFVIII 349 DLYSDMDVVRDDESSPFIQIRSVAKKHPTVWVHYIAEEDWDYAP
PIGFVIII 349 DLYSDMDVVRDDESSPFIQIRSVAKKHPTVWVHYIAEEDWDYAP
MURFVIII 351 DLYSDMDVVRDDESSPFIQIRSVAKKHPTVWVHYIAEEDWDYAP
CANFVIII 343 DLYSDMDVVRDDESSPFIQIRSVAKKHPTVWVHYIAEEDWDYAP

HUMFVIII 399 VPPDPRSYSKSYLNGGPPRIGRKYKKVRFMAYTDETFKTREALCHESGI
PIGFVIII 399 VPPDPRSYSKSYLNGGPPRIGRKYKKVRFVAYTDETFKTREALCHESGI
MURFVIII 399 VPPDPRSYSKSYLNGGPPRIGRKYKKVRFIAYTDETFKTREALCHESGI
CANFVIII 393 VPPDPRSYSKSYLNGGPPRIGRKYKKVRFVAYTDETFKTREALCHESGI

HUMFVIII 449 LGPLLTGEVGDITLLIFKNQASRPYNIYPHGITDVSPHLHRLPKGVKHL
PIGFVIII 449 LGPLLTGEVGDITLLIFKNQASRPYNIYPHGITDVSPHLHRLPKGVKHL
MURFVIII 449 LGPLLTGEVGDITLLIFKNQASRPYNIYPHGITDVSPHLHRLPKGIKHV
CANFVIII 443 LGPLLTGEVGDITLLIFKNQASRPYNIYPHGITDVSPHLHRLPKGVKHL

HUMFVIII 499 KDLPILPGEIFKYKWTVTVEGDPTKSDPRCLTRYSSSFINERDLASGLI
PIGFVIII 499 KDMPILPGEIFKYKWTVTVEGDPTKSDPRCLTRYSSSFINERDLASGLI
MURFVIII 499 KDLPILPGEIFKYKWTVTVEGDPTKSDPRCLTRYSSSFINERDLASGLI
CANFVIII 493 KDMPILPGEIFKYKWTVTVEGDPTKSDPRCLTRYSSSFINERDLASGLI

HUMFVIII 549 GPLLICYSKESVDQRGNQMSDKRNWILFSVFEDENSWYLTENIQRFELNP
PIGFVIII 549 GPLLICYSKESVDQRGNQMSDKRNWILFSVFEDENSWYLTENIQRFELNP
MURFVIII 549 GPLLICYSKESVDQRGNQMSDKRNWILFSVFEDENSWYLTENIQRFELNP
CANFVIII 543 GPLLICYSKESVDQRGNQMSDKRNWILFSVFEDENSWYLTENIQRFELNP

```

HUMFVIII 590 ~~AGVQ~~LDPEFQASNMHSINGTVFEDSLQLSVCLHEVAYWILSIGAQTDF
 PIGFVIII 590 ~~AGVQ~~LDPEFQASNMHSINGTVFEDSLQLSVCLHEVAYWILSIGAQTDF
 MURFVIII 590 ~~AGVQ~~LDPEFQASNMHSINGTVFEDSLQLSVCLHEVAYWILSIGAQTDF
 CANFVIII 593 ~~AGVQ~~LDPEFQASNMHSINGTVFEDSLQLSVCLHEVAYWILSIGAQTDF

HUMFVIII 649 LSVFFSGYTFKHKMVIYEDTLTLFPFSGETVFMSENPNGLWLGLGCHNSDFR
 PIGFVIII 649 LSVFFSGYTFKHKMVIYEDTLTLFPFSGETVFMSENPNGLWLGLGCHNSDFR
 MURFVIII 649 LSVFFSGYTFKHKMVIYEDTLTLFPFSGETVFMSENPNGLWLGLGCHNSDFR
 CANFVIII 643 LSVFFSGYTFKHKMVIYEDTLTLFPFSGETVFMSENPNGLWLGLGCHNSDFR

HUMFVIII 699 NRGMTALLKYSSCDENNGDYEDSYHDIEATLLNNNNVIEPRSFQNSRH
 PIGFVIII 699 NRGMTALLKYSSCDENNGDYEDSYHDIEATLLNNNNVIEPRSFQNSRH
 MURFVIII 699 NRGMTALLKYSSCDENNGDYEDSYHDIEATLLNNNNVIEPRSFQNSRH
 CANFVIII 693 NRGMTALLKYSSCDENNGDYEDSYHDIEATLLNNNNVIEPRSFQNSRH

HUMFVIII 749 PSTRKQKSFATTIPENDIEKIDPQSGERTQCLNQSVDLMLLQCGH
 PIGFVIII 749 PSTRKQKSFATTIPENDIEKIDPQSGERTQCLNQSVDLMLLQCGH
 MURFVIII 749 PSTRKQKSFATTIPENDIEKIDPQSGERTQCLNQSVDLMLLQCGH
 CANFVIII 743 PSTRKQKSFATTIPENDIEKIDPQSGERTQCLNQSVDLMLLQCGH

HUMFVIII 798 PTPHGLLSDLOEAKYESTDDSPGATNNSSSSSSTPQLHHSNDR
 PIGFVIII 797 PTPHGLLSDLOEAKYESTDDSPGATNNSSSSSSTPQLHHSNDR
 MURFVIII 799 PTPHGLLSDLOEAKYESTDDSPGATNNSSSSSSTPQLHHSNDR
 CANFVIII 792 PTPHGLLSDLOEAKYESTDDSPGATNNSSSSSSTPQLHHSNDR

HUMFVIII 848 VFTPEQLQLRLNELGTTVELKKLDKVVSTSNNLSTIPSDNLA
 PIGFVIII 845 VFTPEQLQLRLNELGTTVELKKLDKVVSTSNNLSTIPSDNLA
 MURFVIII 849 VFTPEQLQLRLNELGTTVELKKLDKVVSTSNNLSTIPSDNLA
 CANFVIII 840 VFTPEQLQLRLNELGTTVELKKLDKVVSTSNNLSTIPSDNLA

HUMFVIII 895 AGTDNTSLGPPNPVYHSLGTLTDFGNSSSLTSCDPLSEBNSDS
 PIGFVIII 881 AGTDNTSLGPPNPVYHSLGTLTDFGNSSSLTSCDPLSEBNSDS
 MURFVIII 898 AATEKTSLGPPNPVYHSLGTLTDFGNSSSLTSCDPLSEBNSDS
 CANFVIII 890 AATEKTSLGPPNPVYHSLGTLTDFGNSSSLTSCDPLSEBNSDS

HUMFVIII 946 KLESGLMNSQSSSLGNVYSSSRLEFRAHGPALTKDNALFKVNI
 PIGFVIII 928 KLESGLMNSQSSSLGNVYSSSRLEFRAHGPALTKDNALFKVNI
 MURFVIII 948 WILDSGLMNSQSSSLGNVYSSSRLEFRAHGPALTKDNALFKVNI
 CANFVIII 940 KLESGLMNSQSSSLGNVYSSSRLEFRAHGPALTKDNALFKVNI

HUMFVIII 996 SLKTNFATNATNRKTHIDPELLIENSTSVWQNIESTEFKEVTS
 PIGFVIII 968 SLKTNFATNATNRKTHIDPELLIENSTSVWQNIESTEFKEVTS
 MURFVIII 998 SLKTNFATNATNRKTHIDPELLIENSTSVWQNIESTEFKEVTS
 CANFVIII 990 SLKTNFATNATNRKTHIDPELLIENSTSVWQNIESTEFKEVTS

HUMFVIII 1045 LIHNTMDKNITALLNHNMSNKTITSSKNHNTGKKFPIPIADANPDM
 PIGFVIII 1001 LIHNTMDKNITALLNHNMSNKTITSSKNHNTGKKFPIPIADANPDM
 MURFVIII 1047 LIHNTMDKNITALLNHNMSNKTITSSKNHNTGKKFPIPIADANPDM
 CANFVIII 1040 LIHNTMDKNITALLNHNMSNKTITSSKNHNTGKKFPIPIADANPDM

HUMFVIII 1095 SEFKMLFELPESNMIRTHGKNSLSEQSPSPKQLSLGSEKSVKQNFEL
 PIGFVIII 1048 SEFKMLFELPESNMIRTHGKNSLSEQSPSPKQLSLGSEKSVKQNFEL
 MURFVIII 1097 SEFKMLFELPESNMIRTHGKNSLSEQSPSPKQLSLGSEKSVKQNFEL
 CANFVIII 1090 SEFKMLFELPESNMIRTHGKNSLSEQSPSPKQLSLGSEKSVKQNFEL

HUMFVIII 1145 SEKNKVVVSEDBFTKDYOLKEMYFSEKRLFLTNLNNNNHNNHCKEKT
 PIGFVIII 1085 SEKNKVVVSEDBFTKDYOLKEMYFSEKRLFLTNLNNNNHNNHCKEKT
 MURFVIII 1147 SEKNKVVVSEDBFTKDYOLKEMYFSEKRLFLTNLNNNNHNNHCKEKT
 CANFVIII 1136 SEKNKVVVSEDBFTKDYOLKEMYFSEKRLFLTNLNNNNHNNHCKEKT


```

HUMFVIII 1195 QEEIEKKEQLIGENVVLPQIHITGTKNFMKNLFLSTRONVEQSTQCAI
PIGFVIII 1102 SEMERREKLVQENVLPQVITATGKNELRNIFEGSTEDSVEGETQSH
MURFVIII 1197 QEEIEKKEQLIEENVVLPQVHITATGSKNFLKDIILSTRONVEQSTQCAI
CANFVIII 1184 QEEIERKEKLQENVVLPQSHITGTKNFLKNLFLSTRONVEQSTQCAI

HUMFVIII 1245 APVLQDERSLNDSTNNTNIAHFSKQSEANLEGLGNQTKQIVEYR
PIGFVIII 1152 APVLQDERSLNDSEANLHAFSRISEANLEAGNRTESEYR
MURFVIII 1244 APVLQDERSLNDSTNNTNIAHFSKQSEANLEGLGNQTKQIVEYR
CANFVIII 1234 APVLQDERSLNDSEANLHAFSRISEANLEGLGNQTKQIVEYR

```

HUMFVII 1293 STPRILNLSGQVITQSRKRLKQRLPLEEISERKRIIVDDTSTQWSK
 PIGFVII 1192SPLEELISERKRIIVDDTSTQWSK
 MURFVII 1293PSQKRIITQSRKRLKQRLPLEEISERKRIIVDDTSTQWSK
 CANFVII 1282 STPRILNLSGQVITQSRKRLKQRLPLEEISERKRIIVDDTSTQWSK

HUMFVII 1343 NNKLLSTPLTQIDYNEKEKRAITQSPLSDCNRRNRRIRNNSSLPPIAK
 PIGFVII 1234KRAITQSPLSDCNRRNRRIRNNSSLPPIAK
 MURFVII 1319 NNKLLSTPLTQIDYNEKEKRAITQSPLSDCNRRNRRIRNNSSLPPIAK
 CANFVII 1331 NNKLLSTPLTQIDYNEKEKRAITQSPLSDCNRRNRRIRNNSSLPPIAK

HUMFVII 1393 NSSPFSIRDIILTRVTPQNNSSHLPAAS...YKRRPSGVQESSHFLOEA
 PIGFVII 1234NSSPFSIRDIILTRVTPQNNSSHLPAAS...YKRRPSGVQESSHFLOEA
 MURFVII 1368 NSSPFSIRDIILTRVTPQNNSSHLPAAS...YKRRPSGVQESSHFLOEA
 CANFVII 1381 NSSPFSIRDIILTRVTPQNNSSHLPAAS...YKRRPSGVQESSHFLOEA

HUMFVII 1439 KNNLSLALTLTLEMTQGGREVSLSGTSATNSVTYKKVENTVILKPLLEET
 PIGFVII 1243 KNNLSLALTLTLEMTQGGREVSLSGTSATNSVTYKKVENTVILKPLLEET
 MURFVII 1415 KNNLSLALTLTLEMTQGGREVSLSGTSATNSVTYKKVENTVILKPLLEET
 CANFVII 1431 KNNLSLALTLTLEMTQGGREVSLSGTSATNSVTYKKVENTVILKPLLEET

HUMFVII 1489 SGKVELLPSVHYHVEDIIPPTSTNSGSPGHLDLVPSLSLQTEGKIKWNEA
 PIGFVII 1293 SGKVELLPSVHYHVEDIIPPTSTNSGSPGHLDLVPSLSLQTEGKIKWNEA
 MURFVII 1465 SGKVELLPSVHYHVEDIIPPTSTNSGSPGHLDLVPSLSLQTEGKIKWNEA
 CANFVII 1481 SGKVELLPSVHYHVEDIIPPTSTNSGSPGHLDLVPSLSLQTEGKIKWNEA

HUMFVII 1539 NRPGKVPPFLKATESSKTPSKLLPLAWDNHYETIIPKEEWSQEKSP
 PIGFVII 1343 NRPGKVPPFLKATESSKTPSKLLPLAWDNHYETIIPKEEWSQEKSP
 MURFVII 1515 NRPGKVPPFLKATESSKTPSKLLPLAWDNHYETIIPKEEWSQEKSP
 CANFVII 1531 NRPGKVPPFLKATESSKTPSKLLPLAWDNHYETIIPKEEWSQEKSP

HUMFVII 1589 STAFKKKDDTIILSLNNKSNHAIAAINEGQNKPEIETWAKQGRTERLCS
 PIGFVII 1370 STAFKKKDDTIILSLNNKSNHAIAAINEGQNKPEIETWAKQGRTERLCS
 MURFVII 1563 STAFKKKDDTIILSLNNKSNHAIAAINEGQNKPEIETWAKQGRTERLCS
 CANFVII 1581 STAFKKKDDTIILSLNNKSNHAIAAINEGQNKPEIETWAKQGRTERLCS

HUMFVII 1638 QNPVPLKRHQREITVTTTQSGGSDYDDTISVEKKEDFDIYEDENQS
 PIGFVII 1420 QNPVPLKRHQREITVTTTQSGGSDYDDTISVEKKEDFDIYEDENQS
 MURFVII 1611 QNPVPLKRHQREITVTTTQSGGSDYDDTISVEKKEDFDIYEDENQS
 CANFVII 1630 QNPVPLKRHQREITVTTTQSGGSDYDDTISVEKKEDFDIYEDENQS

HUMFVII 1688 PRSPQKKTRHYFIAAVERLWDYGMSSSFHVLNRRAQSGVPPQFKKVVFP
 PIGFVII 1470 PRSPQKKTRHYFIAAVERLWDYGMSSSFHVLNRRAQSGVPPQFKKVVFP
 MURFVII 1658 PRSPQKKTRHYFIAAVERLWDYGMSSSFHVLNRRAQSGVPPQFKKVVFP
 CANFVII 1680 PRSPQKKTRHYFIAAVERLWDYGMSSSFHVLNRRAQSGVPPQFKKVVFP

HUMFVII 1738 FTDGSPFTQPLTYRGE LNEHLGLLGPTIRAEVEDNIMVTFRNQASRPYSFYS
 PIGFVII 1520 FTDGSPFTQPLTYRGE LNEHLGLLGPTIRAEVEDNIMVTFRNQASRPYSFYS
 MURFVII 1707 FTDGSPFTQPLTYRGE LNEHLGLLGPTIRAEVEDNIMVTFRNQASRPYSFYS
 CANFVII 1730 FTDGSPFTQPLTYRGE LNEHLGLLGPTIRAEVEDNIMVTFRNQASRPYSFYS

HUMFVII 1788 SLISYEEDQSGAEPRKNPVKPNETKTYFWKVQHMAPTDEFDCKAWAY
 PIGFVII 1570 SLISYEEDQSGAEPRKNPVKPNETKTYFWKVQHMAPTDEFDCKAWAY
 MURFVII 1757 SLISYEEDQSGAEPRKNPVKPNETKTYFWKVQHMAPTDEFDCKAWAY
 CANFVII 1780 SLISYEEDQSGAEPRKNPVKPNETKTYFWKVQHMAPTDEFDCKAWAY

HUMFVII 1838 FSDVDLEKDVHSGLIGPLLICANTLNPAHGRQVTVQEFALPFTI FDETK
 PIGFVII 1620 FSDVDLEKDVHSGLIGPLLICANTLNPAHGRQVTVQEFALPFTI FDETK
 MURFVII 1806 FSDVDLEKDVHSGLIGPLLICANTLNPAHGRQVTVQEFALPFTI FDETK
 CANFVII 1830 FSDVDLEKDVHSGLIGPLLICANTLNPAHGRQVTVQEFALPFTI FDETK

HUMFVIIII 1888 SWYPTENMERNCRAPONTMEDPTKENTRPHAINGYIMDTLPGLVMAQD
 PIGFVIIII 1870 SWYPTENVERNCRAPONTMEDPTLKENYRPHAINGYVMDTLPGLVMAQD
 MURFVIIII 1856 SWYPTENVERNCRAPONTMEDPTLKENYRPHAINGYVMDTLPGLVMAQD
 CANFVIIII 1880 SWYPTENLERNCRAPONTMEDPTLKENYRPHAINGYVMDTLPGLVMAQD

HUMFVIIII 1938 QRIRWYLLSMGSGENIHSIHFSCHVFTVRKKEEYKMAVYNLYPGVFETVE
 PIGFVIIII 1720 QRIRWYLLSMGSGENIHSIHFSCHVFTVRKKEEYKMAVYNLYPGVFETVE
 MURFVIIII 1906 QRIRWYLLSMGSGENIHSIHFSCHVFTVRKKEEYKMAVYNLYPGVFETVE
 CANFVIIII 1930 QRIRWYLLSMGSGENIHSIHFSCHVFTVRKKEEYKMAVYNLYPGVFETVE

HUMFVIIII 1988 MLPSKQGIWRVECLIGEHLQAGMSTLPLVYSKQQTPLGMA SGHIRDPQI
 PIGFVIIII 1770 MLPSKQGIWRVECLIGEHLQAGMSTLPLVYSKQQTPLGMA SGHIRDPQI
 MURFVIIII 1956 MLPSKQGIWRVECLIGEHLQAGMSTLPLVYSKQQTPLGMA SGHIRDPQI
 CANFVIIII 1980 MLPSKQGIWRVECLIGEHLQAGMSTLPLVYSKQQTPLGMA SGHIRDPQI

HUMFVIIII 2038 TASGQYQGQWAPKLARLHYSGSINAWSTKDPFWSWKVDLLAPMIHGIQTQ
 PIGFVIIII 1820 TASGQYQGQWAPKLARLHYSGSINAWSTKDPFWSWKVDLLAPMIHGIQTQ
 MURFVIIII 2006 TASGQYQGQWAPKLARLHYSGSINAWSTKDPFWSWKVDLLAPMIHGIQTQ
 CANFVIIII 2030 TASGQYQGQWAPKLARLHYSGSINAWSTKDPFWSWKVDLLAPMIHGIQTQ

HUMFVIIII 2088 GARQKFSSLYISQFIIMYSLDGKKWQSYTRGNSGTGLMVFFGNVDSGQIKH
 PIGFVIIII 1870 GARQKFSSLYISQFIIMYSLDGKKWQSYTRGNSGTGLMVFFGNVDSGQIKH
 MURFVIIII 2056 GARQKFSSLYISQFIIMYSLDGKKWQSYTRGNSGTGLMVFFGNVDSGQIKH
 CANFVIIII 2080 GARQKFSSLYISQFIIMYSLDGKKWQSYTRGNSGTGLMVFFGNVDSGQIKH

HUMFVIIII 2138 NIFNPPPIIARYIRLHPHTYSIRSTLRMLMGCDLNSCSMPLGMESKAISD
 PIGFVIIII 1920 NIFNPPPIIARYIRLHPHTYSIRSTLRMLMGCDLNSCSMPLGMESKAISD
 MURFVIIII 2106 NIFNPPPIIARYIRLHPHTYSIRSTLRMLMGCDLNSCSMPLGMESKAISD
 CANFVIIII 2130 NIFNPPPIIARYIRLHPHTYSIRSTLRMLMGCDLNSCSMPLGMESKAISD

HUMFVIIII 2188 AQITASSYPTNMFAWSPSQARLHLQGRTHNAWRPQYNNPKEWLQVDQKT
 PIGFVIIII 1970 AQITASSYPTNMFAWSPSQARLHLQGRTHNAWRPQYNNPKEWLQVDQKT
 MURFVIIII 2156 AQITASSYPTNMFAWSPSQARLHLQGRTHNAWRPQYNNPKEWLQVDQKT
 CANFVIIII 2180 AQITASSYPTNMFAWSPSQARLHLQGRTHNAWRPQYNNPKEWLQVDQKT

HUMFVIIII 2238 MKVTOITQGVKSLTSMYVKEFLISSSQDGHWTFLQNGKVKVFQGNQ
 PIGFVIIII 2020 MKVTOITQGVKSLTSMYVKEFLISSSQDGHWTFLQNGKVKVFQGNQ
 MURFVIIII 2206 MKVTOITQGVKSLTSMYVKEFLISSSQDGHWTFLQNGKVKVFQGNQ
 CANFVIIII 2230 MKVTOITQGVKSLTSMYVKEFLISSSQDGHWTFLQNGKVKVFQGNQ

HUMFVIIII 2288 DSSTPVVNSLDPPLLTRYLRIRHPQSWAHIALRLEVLGCEAQCLY
 PIGFVIIII 2070 DSSTPVVNSLDPPLLTRYLRIRHPQSWAHIALRLEVLGCEAQCLY
 MURFVIIII 2256 DSSTPVVNSLDPPLLTRYLRIRHPQSWAHIALRLEVLGCEAQCLY
 CANFVIIII 2280 DSSTPVVNSLDPPLLTRYLRIRHPQSWAHIALRLEVLGCEAQCLY

Exhibit 2: Lundblad et al., *Thromb. Haemost.* 84:942-948 (2000)

Review Article

Issues with the Assay of Factor VIII Activity in Plasma and Factor VIII Concentrates

Roger L. Lundblad^{1,2*}, Henry S. Kingdon³, Kenneth G. Mann⁴, Gilbert C. White⁵From the ¹Department of Pathology, University of North Carolina, Chapel Hill,²Baxter Healthcare – Hyland Immuno Division, Duarte, CA,³Baxter Healthcare – Hyland Immuno Division (Lake Cook, IL),⁴Department of Biochemistry, University of Vermont, Burlington, VT and the⁵Department of Medicine, Division of Hematology-Oncology, University of North Carolina at Chapel Hill, USA

Key words

Factor VIII, factor VIII assay/concentrates, chromogenic assays

Summary

A review of the literature suggests that assays accurate for the determination of factor VIII in plasma samples may not necessarily retain this accuracy when used for the determination of factor VIII in high-purity factor VIII concentrates such as Hemofil[®] M. Review of assay data suggests that it is imperative to obtain maximal activation of the factor VIII in the sample with thrombin when using an assay system of isolated coagulation factors such as the two-stage assay or the various chromogenic substrate assays. Based on a combination of ease and reproducibility of performance and correlation of *in vivo* and *in vitro* measurements, it is recommended that the one-stage activated partial thromboplastin time performed with plasma from an individual with severe hemophilia A be used for the measurement of factor VIII potency. Chromogenic substrate assays can be used if care is taken to assure optimal activation of factor VIII by thrombin in the assay and the presence of sufficient factor IXa, phospholipid and calcium ions to stabilize factor VIIIa during the assay process.

Introduction

The accurate assay of factor VIII is critical in the management of hemophilia A and will be of increasing importance as new modes of therapy such as continuous infusion and other prophylactic approaches are developed (1-5). A desirable criterion for such an assay is that it should be equally sensitive and accurate with therapeutic concentrates and patient samples after infusion. It is equally important for manufacturers of therapeutic concentrates to have a robust and accurate assay for use in the assignment of potency. The World Health Organization (WHO) and the International Society on Thrombosis and Haemostasis (ISTH) have long been involved in calibrating and distributing standards

for use in these assays (6, 7). It has been understood that a concentrate standard (e.g. MEGA I) must be used when assigning concentrate potency (8, 9) and that a plasma standard be used when calibrating clinical assays although there can be issues with this approach (10). It is important, however, that concentrate and plasma assay correlate well in order to make sense out of the clinical results.

The activated partial thromboplastin time (aPTT) (11, 12) has been used extensively for the factor VIII assay both in therapeutic concentrates and in hemophilia A patients. There has been concern that the aPTT does not provide an accurate measurement of the potency of highly purified factor VIII (13, 14). It has been suggested that differences between values obtained with the aPTT and a chromogenic or two-stage assay reflects the presence of activated factor VIII in these preparations (15-17). Another emerging complication for factor VIII assays is the European requirement for the use of chromogenic substrate assay systems for factor VIII (18). The European Pharmacopoeia describes a generic chromogenic substrate assay and gives recommendations on the level of purity of the factors used to formulate the assays (generally around 50%) and the concentration of the various components. The method recommends that the reaction to activate the factor X be terminated before all the factor X is activated. The package inserts of the various commercial chromogenic substrate assays do not always contain enough information to know if they meet the criteria in the Pharmacopoeia.

The purpose of this review is to critically examine the various assay systems currently available for the assay of factor VIII activity both in therapeutic concentrates and plasma samples. The review also considers the advances in the understanding of the biochemistry of factor VIII that are relevant to the various assay systems. The reader is referred to several recent reviews of factor VIII for general information (19-24).

Mechanism of Action of Factor VIII in Blood Coagulation

A major problem in the assay of factor VIII is that this protein can only be measured in an indirect manner. Factor VIII is not an enzyme but rather a procofactor in the activation of factor X by factor IXa (19-24). Thus, all currently available factor VIII assays measure factor VIII indirectly through its cofactor activity (available only after peptide bond cleavage) to generate factor Xa. Factor Xa is measured

Correspondence to: Dr. Roger L. Lundblad, P.O. Box 16695, Chapel Hill, NC 27516-6695, USA. Tel.: +01 919 960 8889; Fax: +01 919 960 8889; E-mail: r.lundblad@worldnet.att.net

either in a direct manner such as described below for the chromogenic substrate assay systems or in an indirect manner through the formation of fibrin by thrombin formed by the coagulation cascade.

Activation State of Factor VIII

The assay of factor VIII would be less complicated if factor VIII did not require activation prior to function as a cofactor in the activation of factor X (25, 26). Activation of factor VIII by thrombin results in an observed 30–50 fold (27, 28) or greater (29) increase in cofactor activity. However, the full extent of potential factor VIII activation is not known since the activity of single-chain factor VIII, unlike that of the homologous molecule, factor V (30), has not been measured. In the absence of specific data to the contrary, it is generally assumed that single chain factor VIII does not have any cofactor activity (31, 32). However, estimates of baseline activity are flawed also because as with factor VIII, all assays are indirect. Understanding of the activation process is further complicated because species of factor VIII which have undergone some, but not all of the peptide bond cleavages required for full activation, demonstrate enhanced cofactor activity corresponding to the degree of processing (33). For example, it was determined that cleavage of the 80 KD light chain to the 70 KD $A_2C_2C_3$ fragment is associated with a six-fold increase in activity (33). Cleavage of the heavy chain at residues 372–373 yields the heterotrimer, $A_2A_2A_2C_2C_3$ (34, 35) and resulted in additional activation but the product was quite unstable. Recent studies (36, 37) with abnormal factor VIII proteins isolated from patients with mild hemophilia A have provided additional evidence for the importance of light chain cleavage at 1689–1690 during the activation process. These studies demonstrated that abnormal factor VIII proteins cleaved poorly at residues 372–373, are still partially activated by thrombin via light chain cleavage. Previously, this cleavage was considered to be not as critical as heavy chain cleavage for the activation process (38, 39); the change in perspective occurred when it became appreciated that light chain cleavage is required for von Willebrand factor (vWF) dissociation and vWF dissociation from factor VIII is required for cofactor activity (39). The importance of light chain cleavage in the factor VIII activation process has been further demonstrated by Regan and Fay (40) in experiments where the isolated thrombin-cleaved light chain was allowed to associate with the isolated heavy chain. The activity of this hybrid protein was some three-fold greater than the factor VIII obtained from the association of intact light chain and heavy chain.

Proteases other than thrombin have been demonstrated to activate factor VIII. Factor Xa is the most likely protease to augment the activity of thrombin. While factor Xa can partially activate factor VIII, it is clear that complete activation is accomplished only with thrombin (41, 42). It has been reported that factor IXa can activate Factor VIII but the contribution of this reaction to factor VIII cofactor function would be far less significant than that catalyzed by thrombin or factor Xa (43). In vitro studies would suggest that thrombin activation is requisite for biological function (25, 27, 44–50). It is emphasized that factor VIII, once activated to factor VIIIa, decays quite rapidly.

Influence of von Willebrand Factor on Factor VIII Activity

The presence of von Willebrand Factor (vWF) appears to influence the in vitro assay of factor VIII. Highly purified recombinant factor VIII preparations which do not contain vWF gave lower-than-expected potency values when assayed in vWF-depleted factor VIII-deficient plasmas (51, 52). As a result, dilution of highly-purified factor VIII

samples in hemophilia A plasma prior to assay has been recommended (53–55). In support of this concept, it has been shown that plasma vWF is required for a normal circulating half-life for factor VIII (56) and that vWF-free recombinant factor VIII associates with vWF after infusion (57). In addition to this effect on the state of factor VIII in the circulation, vWF also directly stabilized factor VIII (58).

While the results with the use of vWF-antibody-immunodepleted factor VIII-deficient plasma may result in an unsatisfactory assay for high-purity concentrates, such substrate plasmas continued to be useful for factor VIII assay in plasma samples (59). Equivalence as a diluent has been reported for hemophilia A plasma and factor VIII-deficient plasma obtained by treatment with EDTA (55). Rothschild et al. (60) have reported that a plasma depleted by both a factor VIII antibody and a vWF antibody yielded equivalent results compared to hemophilia A plasma with intermediate and high purity plasma-derived concentrates.

Clear biochemical evidence for the importance of vWF in a factor VIII assay system has not been demonstrated. vWF does not appear to have an effect on the thrombin-activation of factor VIII with respect to the increase in biological activity but does serve as a cofactor for a non-rate limiting step in the overall activation process (61, 62). These studies were performed with porcine factor VIII and it is not clear that such results would be obtained with the human protein. However, if these results could be effectively translated to the human system, a role for vWF in the assay systems would be clear considering the essential role of thrombin in the factor VIII activation process. For example, Vukovich et al. (63) presented data supporting a role for vWF in a two-stage assay but not in a one-stage assay. Somewhat the opposite conclusion was reached by other investigators who reported that factor VIII with greatly reduced vWF content appeared to have higher activity in the two-stage assay than in the one-stage assay (64).

vWF does appear to inhibit the activation of factor VIII by factor Xa (65, 66). In other studies, vWF has been reported to decrease the stability of both thrombin-modified and factor Xa-modified factor VIII (67). The issue of the influence of vWF on the assay of factor VIII in high purity concentrates versus plasma factor VIII or low-purity concentrates is complicated by the marked differences in vWF quality in the high-purity factor VIII concentrates (68–70). Aronson and Chang showed that while a recombinant factor VIII (Kogenate[®]) and a highly purified factor VIII concentrate obtained by immunoaffinity chromatography with a factor VIII antibody (Hemofili[®] M) readily associated in vitro with von Willebrand factor to form a more-rapidly sedimenting complex on ultracentrifugal analysis (71), another highly-purified concentrate obtained by immunoaffinity chromatography with a vWF antibody (Monoclate[®]) did not show this association in vitro with von Willebrand factor. There was no difference in the sedimentation behavior of the three therapeutic preparations after infusion into a hemophilic patient. Differences between Monoclate[®] and Hemofili[®] M have been demonstrated in a chromogenic substrate assay system (14) and in data shown below. The molecular basis for the assay discrepancies is not clear but it is unlikely to be related to the activation state of factor VIII since factor VIIIa (activated factor VIII) does not effectively bind to von Willebrand factor (72).

In Vitro Assay Systems for Factor VIII

Three types of assay systems exist for the measurement of factor VIII in biological samples (73). These are the one-stage partial thromboplastin time using factor VIII-deficient plasma (11, 12), the two-stage assay (74, 75) and chromogenic substrate assays (76, 77).

The one-stage assay for factor VIII measures the ability of a sample to shorten the clotting time of severe hemophilia A plasma after contact activation and recalcification. This assay uses the intrinsic system in which the reaction is initiated via contact activation with fibrinogen clotting as the endpoint. In the one-stage assay for factor VIII, factor VIII-deficient plasma is preincubated with phospholipid containing contact activation material such as kaolin or elagic acid and the sample; the assay itself is initiated by the addition of calcium chloride. The use of this assay assumes that factor VIII in the sample is rate-limiting with all other necessary components (clotting factors, phospholipid or activated platelets and other cofactors) present at saturating levels. The aPTT, while likely the assay system most closely related to physiologic function, has some major drawbacks. The assumption that all components other than factor VIII are present at saturating levels is not necessarily valid and the sample may contain components other than factor VIII which could contribute to or partially oppose the measured activity. An example of this is provided by the observations on the apparent influence of vWF on observed factor VIII activity (51, 52). The presence of trace amounts of factor VIII in the substrate plasma also can influence results (78). Finally, there is evidence that each factor VIII-deficient plasma will give a somewhat unique slope for the standard curve.

The two-stage assay system consists of two separate reaction periods. The first incubation contains the factor VIII sample in combination with a serum reagent which contains human serum (source of factor IXa and factor X), bovine serum absorbed with barium salts (source of factor V/Va), calcium ions and phospholipid. The prothrombinase (79) which is formed is measured in the second stage by the ability to activate prothrombin to thrombin with the rate of fibrinogen conversion to fibrin being measured. Since the two-stage assay is an isolated system, there is (a) no physiologically-significant thrombin feedback activation of factor VIII and (b) no modulatory effect of plasma protease inhibitors (serpins).

The third type of assay system available is the chromogenic substrate assay(s). It should be noted that the chromogenic substrate assay systems are so-named not because of a similarity in reaction mechanism, but rather that a colored product, p-nitroaniline, is the signal measured (80). Chromogenic substrate assays are related to the two-stage assay in that there are usually two incubation periods. The first, which actually measures factor VIII cofactor activity, involves the reaction of factor IXa, calcium ions, factor X, phospholipid and the sample or reaction blank; factor Xa is produced during this incubation period. The second incubation period measures the factor Xa produced in the first incubation period, using a peptide nitroanilide substrate. The hydrolysis of the peptide nitroanilide substrate forms p-nitroaniline which is usually measured by absorbance at 405 nm.

Currently, there are four chromogenic substrate assay systems which are commercially available; these chromogenic substrate assays are not equal with respect to mechanism:

- (1) The original Chromogenix system contains factor X, factor IXa, prothrombin, calcium ions, phospholipid and the factor VIII sample/standard/blank.
- (2) The Dade-Behring, the new Chromogenix (81) and Biopool systems contain factor X, factor IXa, thrombin, calcium ions, phospholipid and the factor VIII sample/standard/blank. These three systems contain thrombin which is added to maximize factor VIII activation such that there would be improved molecular homogeneity of factor VIII during assay.

All of these assay systems consist of two stages. The first stage generates factor Xa, the difference between the two types of systems is that the Dade-Behring, the new Chromogenix and the Biopool systems

Table 1 Comparison of two ultra-high purity factor concentrates with two chromogenic assays and the activated partial thromboplastin time

| Sample ^a | aPTT | Dade-Behring | Chromogenix | Label Potency |
|---------------------|-------------------|--------------|-------------|---------------|
| Concentrate A | 21.0 ^b | 20.4 | 17.1 | 23.5 |
| Concentrate B | 98.5 | 86.2 | 30.2 | 100 |

a. Samples had been previously dissolved in deionized water and stored at -70°C. The samples were produced in Hemophilia A plasma prior to assay.

b. Values are presented as units of factor VIII C activity/ml, sample obtained with the Mega I standard.

contain thrombin while with the first Chromogenix assay thrombin must be generated during the first stage incubation. The second stage is essentially identical for the various systems in that there is a chromogenic factor Xa substrate, buffer and a thrombin inhibitor. It is the hydrolysis of the chromogenic substrate that is actually measured in the assay. The rate of hydrolysis will vary between methods depending on the chromogenic substrate used. Although all modern chromogenic substrates have reasonable specificity, some cross-reactivity cannot be discounted. All formulations on the market at present contain thrombin inhibitors preventing thrombin interference in the assay. A new chromogenic substrate assay has been reported by Immuno. The Immuno assay uses all human proteins but contains no thrombin (82).

What about comparison of assay results? As has been mentioned above, there have been continuing reports of discrepancies between assay systems. While not wishing to complicate the issue, Table 1 presents a comparison between two ultra-high purity factor VIII concentrates with three assay systems. The point of this comparison is not to argue for the superiority of one concentrate vs. another but rather to emphasize the importance of thrombin in the chromogenic assay system. It is clear that with either concentrate, the aPTT is the most accurate measure when compared to label potency; the Dade-Behring assay system which contains thrombin would appear to be more accurate than the original Chromogenix system.

Concentrates: Problems in the Assay of High Purity Factor VIII

The accurate determination of factor VIII potency in therapeutic concentrates is of critical importance to all participants: manufacturers, treaters and patients. In the final analysis it is the *in vivo* hemostatic effectiveness which is decisive in dosing considerations. What then is the most reliable *in vitro* assay to predict *in vivo* performance? Our thesis is that the one-stage activated partial thromboplastin time (aPTT) is at present the most accurate assay for the determination of factor VIII potency in therapeutic concentrates. This recommendation is based not on theoretical considerations but rather on the observed correlation with clinical performance with the potency of factor VIII preparations (83). However, recognizing that the chromogenic assays provide advantages in terms of more facile integration into laboratory information management systems and the lack of necessity of using hemophilia A plasma, we submit that a chromogenic assay system could be effectively used for potency assessment and patient management only if there is clinical experience to verify that vial potency designation is meaningful in the determination of dose and that this dosage translates by determination of patient plasma factor VIII activity into satisfactory control of bleeding under the full spectrum of clinical experience.

Factor VIII Assay Results: In Vivo Studies

Earlier studies with less highly-purified concentrates reported that two-stage activity was increased as compared to one-stage activity. When the two assays were used to follow the *in vivo* pharmacokinetics of factor VIII, the one-stage assay gave a yield equivalent to 100% recovery while a lower recovery (80%) was obtained with the two-stage assay (84). Similar results were obtained by other investigators with an intermediate purity concentrate (85). The one-stage assay gave $109 \pm 20\%$ of predicted activity recovery while the two-stage assay gave $92 \pm 14\%$. The Chromogenic Coatest assay gave a similar result ($91 \pm 11\%$). Smaller differences in *in vivo* recovery with intermediate concentrates using the three assay systems were observed by other investigators (86).

While the one-stage assay has been criticized as being unreliable (87), it appears to remain as the assay of choice for the management of patients with hemophilia in the clinic on a global basis (83, 88-100).

Factor VIII Assay Issues to Be Resolved

Specific issues regarding the assay of factor VIII are as follows and must be resolved or reconciled for the successful interpretation of all factor VIII assay results.

First, in any assay system (one-stage, two-stage, chromogenic), factor VIII activity is measured by the rate of factor X activation. Given our current understanding of factor VIII, if the factor VIII sample contains any species other than fully activated (i.e. thrombin-activated) factor VIII, the assay will then, in fact, be a coupled system. With unactivated factor VIII, conversion to factor VIIa must be effectively accomplished before its participation as a cofactor in the activation of factor X; as such, without full activation of factor VIII, it is possible that the assay system would measure the rate of conversion of factor VIII to factor VIIa rather than the amount of factor VIII present. This system is comparable to other coupled assay systems used in clinical chemistry (101-104): in this context, the generation of factor Xa by factor IXa is dependent upon the rate of generation of factor VIIa such that the activation of factor X is coupled to the generation of factor VIIa. Any coupled assay system must be rigorously validated to demonstrate unequivocally that the measured response is directly related to the concentration of the substance being measured. In other words, in the case of factor VIII, the amount of factor Xa formed in a specific time period and measured either by rate of formation (kinetic rate assay) or amount formed after a specific period of time (endpoint assay) must be proportional to the amount of Factor VIII in the system. While this may be accomplished in the chromogenic assays cited herein, it is somewhat more difficult for either the two-stage or one-stage assay systems. With respect to the nature of the factor VIII molecular species present, the activation step must not be rate limiting. Molecular homogeneity would be most easily accomplished in an isolated system (either two-stage or chromogenic) by fully activating the factor VIII sample prior to assay while assuming that there is sufficient factor IXa, calcium ions and phospholipid to stabilize the activated factor VIII during the assay.

Secondly, how does vWF affect factor VIII function and does the quality of the vWF matter? The quality issue refers to whether there is a difference between high-molecular weight vWF, intermediate molecular-weight vWF and vWF dimer. The potential differential effect of vWF (quantitative and qualitative) on the various assay systems must be elucidated for successful interpretation of results; in particular, the influence of the hemophilic plasma used for dilution of the high purity concentrates must be explained.

Third, there is the general issue of the *in vivo* recovery of ultra-high purity factor VIII concentrates with particular emphasis on the recombinant DNA products (Kogenate[®], Recombinate[®] and Refacto[®]) (105). There has been considerable recent interest in problems associated with Refacto[®]. Refacto[®] is a B-domainless recombinant factor VIII therapeutic developed by Pharmacia-Upjohn (106-108) which is being commercialized by Genetics Institute. While primary data is not available with the use of Refacto[®] in hemophilia A subjects, Lusher and colleagues (109) state that "While recovery values are higher with the chromogenic assay than with the one-stage assay for the two full-length recombinant FVIII preparations, a unique discrepancy exists for the B-domainless rFVIII preparation, Refacto[®]. With Refacto[®], recoveries measured by the chromogenic assay are as expected, whereas one-stage methods give values of approximately 50% of that expected." Mikaelsson et al. (110) reported that there are marked differences in the phospholipid requirement in the assay of Refacto[®] compared to the other factor VIII preparations. This might explain the differences observed by Lusher et al. (109). In earlier studies on B-domainless factor VIII, Pittman et al. (111) note similar activity values for either the Coatest chromogenic or the one-stage assay. In a study of some structurally-related factor VIII derivatives, Mertens et al. (112) report large differences between the chromogenic assay (cofactor activity) and the one-stage assay with hemophilia A substrate plasma (procoagulant activity). This is a complex problem which will require study in larger patient populations.

Finally, regardless of the type of assay chemistry being used, in order to have valid assays one must adhere to appropriate statistical standards (c.f. 113). An appropriate dilution curve of both sample and standard must demonstrate linearity, a significant regression (i.e. concentration dependence) and the lines for standard and sample must be parallel.

Conclusions

Given the current status of information on the assay of factor VIII in therapeutic concentrates, we suggest the following recommendation program.

1. The aPTT assay is an accurate and sensitive method for both the determination of factor VIII potency in therapeutic concentrates and for the subsequent measurement of factor VIII in hemophilic recipients.
2. Chromogenic Assays can also be accurate measures of factor VIII activity both in high-purity concentrates and patient samples if there is an appropriate thrombin activation step included in the assay process. Failure to include this step will result in an inaccurate measure of factor VIII activity in high-purity concentrates. Relative to this requirement is the need for the presence of sufficient factor IXa, calcium ions and phospholipid to stabilize factor VIIa during the assay. An equally important requirement is the demonstration of the clinical relevance of the potency designation as established by the chromogenic assay system.

Acknowledgements

The authors wish to thank Drs. Gordon Bray, Edward Gonnerts, Barry Woodham, Jerry Powell, Don Baker and Professor Hans Peter Schwarz for their helpful comments in the preparation of this manuscript.

References

- Hawkins TE, Green GJ, Romeril K, Milcic GS, Carter JM. Treatment of haemophilia A by continuous factor VIII infusion. *Aust NZ J Med* 1995; 25: 37-9.
- Schulman S, Grief S, Martinowitz U. Stability of factor VIII concentrates after reconstitution. *Am J Hematol* 1994; 45: 217-23.
- Hurst D, Zabor S, Malani D, Miller D. Evaluation of recombinant factor VIII (Kogenate®) stability for continuous infusion using a minipump infusion device. *Haemophilia* 1998; 4: 785-9.
- Curie B, Limentani A, Rosenfeld CG. A practical guide to the evaluation and treatment of hemophilia. *Blood* 1994; 84: 3-9.
- Carlsson MS, Stenberg PB, Björnsson KA, Eklund LH. Pharmaceutical services in the treatment of haemophilic patients. *Amer J Hosp Pharm* 1993; 50: 2073-6.
- Bangham DR, Biggs R, Brozovic M, Denison KWE, Skegg JL. A biological standard for measurement of blood coagulation factor VIII activity. *Bull WHO* 1971; 45: 337-51.
- Blombäck M et al. Nomenclature of quantities and units in thrombosis and haemostasis (Recommendation 1993). *Thromb Haemostasis* 1994; 71: 375-94.
- Barrowcliffe TW. Standards for factor VIII. *Scand J Haematol* 1980; 25 (Suppl. 37): 104-9.
- Kirkwood TBL. Problems in the standardization of factor VIII assays. *Scand J Haematol* 1980; 25 (Suppl. 37): 110-5.
- Barrowcliffe TW, Raut S, Hubbard AR. Discrepancies in potency assessment of recombinant factor VIII concentrates. *Haemophilia* 1998; 4: 634-40.
- Langdell RD, Wagner RH, Brinkhous KM. Effect of antihemophilic factor on one-stage clotting tests. A presumptive assay for hemophilia and a simple one-stage antihemophilic factor assay procedure. *J Lab Clin Med* 1953; 41: 637-47.
- Over J. Methodology of the one-stage assay of factor VIII (VIII: C). *Scand J Haematol* 1986; 33 (Suppl. 41): 13-24.
- Dawson NJ, Kemball-Cook G, Barrowcliffe TW. Assay Discrepancies with Highly Purified Factor VIII Concentrates. *Haemostasis* 1989; 19: 131-7.
- Feldman F, Chandra S, Hendriks M, Schreiber AB. Quality Assurance in Production of Plasma Proteins. In: *Quality Assurance in Transfusion Medicine, Vol. II, Methodological Advances and Clinical Aspects*. Rock G, Sejtichian, MJ, eds. Boca Raton, FL: CRC Press 1993; 259-84.
- Barrowcliffe TW, Watton J, Harmon A, Tubbs JF, Kemball-Cook G. Potency of high purity factor VIII concentrates. *Lancet* 1990; ii: 124.
- Kemball-Cook G, Tubbs JE, Dawson NJ, Barrowcliffe TW. The behavior of different factor VIII concentrates in a chromogenic factor X-activating system. *Brit J Haematol* 1993; 84: 273-8.
- Niemetz J, Nossel HL. Activated Coagulation Factors. In: *Vivo and in-vivo Studies*. Brit J Haematol 1969; 16: 337-51.
- Assay of Blood Coagulation Factor VIII. *European Pharmacopoeia, Third Edition*, Strasbourg: Council of Europe 1997, Section 2.7.4 111-3.
- Hoyer LW. The factor VIII complex: Structure and function. *Blood* 1981; 58: 1-13.
- Foster PA, Zimmerman TS. Factor VIII structure and function. *Blood* 1989; 73: 180-91.
- Kane WH, Davie EW. Blood coagulation factors V and VIII: Structure and functional similarities and their relationships to hemorrhagic and thrombotic disorders. *Blood* 1988; 71: 539-55.
- Chavin ST, Fay PJ. The Purification, Structure and Function of Factor VIII. In: *Factor VIII-von Willebrand Factor*. Segelbacher MJ, Savidge GF, eds. Boca Raton, FL: CRC Press 1989; 2-23.
- Fay PJ. Factor VIII Structure and Function. *Thromb Haemostasis* 1993; 70: 63-7.
- Leiting PJ, van Mourik JA, Mertens K. The life cycle of factor VIII in view of its structure and function. *Blood* 1998; 92: 3983-96.
- Rapaport SI, Schiffman S, Patch MJ, Anes SB. The importance of activation of antihemophilic globulin and proaccelerin by traces of thrombin in the generation of intrinsic prothrombinase activity. *Blood* 1963; 21: 221-36.
- Rapaport SI, Hjort PF, Patch MJ. Further evidence that thrombin-activation of factor VIII is an essential step in intrinsic clotting. *Scand J Clin Invest* 1965; Suppl. 84: 88-100.
- Rick ME, Hoyer LW. Activation of low molecular weight fragment of antihemophilic factor (factor VIII) by thrombin. *Nature* 1974; 252: 404-5.
- Rick M, Hoyer LW. Thrombin activation of factor VIII: A comparison of purified factor VIII and the low molecular weight factor VIII procoagulant. *Brit J Haematol* 1978; 38: 107-19.
- van Diegen G, Tans G, Rosing J, Hemker HC. The role of phospholipid and factor VIIIa in the activation of bovine factor X. *J Biol Chem* 1981; 256: 3433-41.
- Krishnaswamy S, Church WE, Nesheim ME, Mann KG. Activation of Human Prothrombin by Human Prothrombinase. Influence of factor Va on the reaction mechanism. *J Biol Chem* 1987; 262: 3291-9.
- Jesty J. Interaction of feedback control and product inhibition in the activation of factor X by factors IXa and VIII. *Haemostasis* 1991; 21: 208-18.
- Neuenschwander P, Jesty J. The use of acetylated factor X to prevent feedback activation of factor VIII during factor X activation: A tool for kinetic studies. *Analyt Biochem* 1990; 184: 347-52.
- Bihoreau N, Sauger A, Yon JM, Van de Pol H. Isolation and characterization of different activated forms of factor VIII, the human antihemophilic A factor. *Europ J Biochem* 1989; 185: 111-8.
- Lollar P, Parker CG. Subunit structure of thrombin-activated porcine factor VIII. *Biochemistry* 1989; 28: 666-74.
- Fay PJ, Hadornis P, Smudzin TM. Human factor VIIIa subunit structure. Reconstitution of factor VIIIa from the isolated A1/A3-C1-C2 dimer and A2 subunit. *J Biol Chem* 1991; 266: 8957-62.
- Johnson DJD, Pemberton S, Acquila M, Mori PG, Tuddenham EGD, O'Brien DP. Factor VIII S373L: Mutation at P1' site confers thrombin cleavage resistance causing mild haemophilia A. *Thromb Haemostasis* 1994; 71 (4): 428-33.
- Arai M, Inabe H, Higuchi M, Antonarakis SE, Kazanietz HH Jr, Fujimaki M, Hoyer LE. Direct characterization of factor VIII in plasma. Detection of a mutation altering a thrombin cleavage site (arginine-372 → histidine). *Proc Natl Acad Sci USA* 1989; 86: 4277-81.
- Pittman DD, Kaufmann RJ. Proteolytic requirements for thrombin activation of antihemophilic factor (Factor VIII). *Proc Natl Acad Sci USA* 1988; 85: 2429-33.
- Hill-Eubanks DC, Parker CG, Lollar P. Differential proteolytic activation of factor VIII-von Willebrand factor complex by thrombin. *Proc Natl Acad Sci USA* 1989; 86: 6508-12.
- Regan LM, Fay PJ. Cleavage of factor VIII high chain is required for maximal generation of factor VIIIa activity. *J Biol Chem* 1995; 270: 8546-52.
- Neuenschwander P, Jesty J. A comparison of phospholipid and platelets in the activation of factor VIII by thrombin and factor Xa and in the activation of factor X. *Blood* 1988; 72: 1761-70.
- Lollar P, Knutson GJ, Fass DJ. Activation of porcine factor VIII: C by thrombin and factor Xa. *Biochemistry* 1985; 24: 8556-64.
- Lamphear BJ, Fay PJ. Proteolytic interactions of factor IXa with human factor VIII and factor VIIIa. *Blood* 1992; 80: 3120-6.
- Switzer MEP, Pizzo SV, McKee PA. Is there a relatively procoagulant-inactive form of normal antihemophilic factor (Factor VIII)? *Blood* 1979; 54: 916-27.
- Ofosu FA, Hirsh J, Esmo CT, Modi GI, Smith LM, Auvari N, Buchanan MR, Fenton JW II, Blajchman MA. Unfractionated heparin inhibits thrombin-catalyzed amplification reactions of coagulation more efficiently than those catalyzed by Factor Xa. *Biochem J* 1989; 257: 143-50.
- Ofosu FA, Sie P, Modi GI, Fernandez F, Buchanan MR, Blajchman MA, Boncu B, Hirsh J. The inhibition of thrombin-dependent positive-feedback reactions is critical to the expression of the anticoagulant effect of heparin. *Biochem J* 1987; 243: 579-88.
- Broden K, Andersson L-O, Sandberg H. Kinetics of activation of human factor VIII by thrombin. *Thromb Res* 1980; 19: 299-307.

48. Pieters J, Lindhout T, Hemphik HC. In situ-generated thrombin is the only enzyme that effectively activates factor VIII and factor V in thromboplastin-activated plasma. *Blood* 1989; 74: 1021-14.
49. Jones KC, Mann KG. A model for the tissue factor pathway to thrombin. II. A mathematical simulation. *J Biol Chem* 1994; 269: 23367-73.
50. Butenas S, van't Veer C, Mann KG. Evaluation of the initiation phase of blood coagulation using ultrasensitive assays for serine proteases. *J Biol Chem* 1997; 272: 21527-33.
51. Barrowcliffe TW, Tubbs JE, Wong MY. Evaluation of factor VIII deficient plasmas. *Thromb Haemost* 1993; 70: 433-7.
52. Lee M, Maglalang EA, Kingdon HS. An effect of predilution on potency assays of factor VIII concentrates. *Thromb Res* 1983; 30: 511-9.
53. Frandsen H, Stender S, Scheibel E. An effect of predilution on VIII:C determination by the one-stage Assay. *Thromb Res* 1987; 48: 195-203.
54. Barrowcliffe TW. Recommendations for the assay of high-purity factor VIII concentrates. *Thromb Haemost* 1993; 70: 876-7.
55. Mazzurri C, Parquet-Gomez A, Goudemand M. Validation of a procedure for potency assessing of a high purity factor VIII concentrate. Comparison of different factor VIII coagulant assays and effect of predilution. *Thromb Haemost* 1990; 64: 251-5.
56. Brinkhous KM, Sandberg H, Garris JB, Mattsson C, Palm M, Griggs T, MS. Purified human factor VIII procoagulant protein: Comparative hemostatic response after infusions into hemophilic and von Willebrand disease dogs. *Proc Natl Acad Sci USA* 1985; 82: 8752-6.
57. Giles AR, Timlin S, Hoogendoorn H, Fournel MA, Ng P, Pancham N. In vivo characterization of recombinant factor VIII in a canine model of hemophilia A (factor VIII deficiency). *Blood* 1988; 72: 335-9.
58. Fass DN, Bowie EJW, Owen CA Jr, Mann KG. Stability of porcine factor VIII. *Thromb Res* 1975; 6: 109-18.
59. Tekse T, Rosblat F, Goodall AH, Kernoff PBA, Middleton S, Chand S, Denison KW, Austen DEG, Tuddeham EGD. Production of factor VIII deficient plasma by immunodepletion using three monoclonal antibodies. *Brit J Haematol* 1987; 66: 497-502.
60. Rothschild C, Amiral J, Adam M, Meyer D. Preparation of factor-VIII-deficient plasma with antibodies and its use for the assay of factor VIII. *Haemost* 1990; 20: 321-8.
61. Hill-Eubanks DC, Lollar P. vWF is a cofactor for thrombin-catalyzed cleavage of the factor VIII light chain. *J Biol Chem* 1990; 265: 17854-8.
62. Lollar P. The association of factor VIII with factor X. *Mayo Clin Proc* 1991; 66: 524-34.
63. Vukovich T, Keller E, Dolschell W. Low-molecular weight factor VIII requires a cofactor for its procoagulant activity. *Thromb Res* 1980; 503-11.
64. Muniean W, Hatfai WE, Montgomery RR. Influence of High-Molecular Weight Factor VIII on the Measurement of Low-Molecular Weight Factor VIII Procoagulant in Different Assay Systems. *Brit J Haematol* 1982; 51: 649-58.
65. Koedam JA, Hamer RJ, Beeser-Visser NH, Bouma BN, Sixma JJ. The effect of vWF on activation of factor VIII by factor Xa. *Europ J Biochem* 1990; 189: 229-34.
66. Koppelman SJ, Koedam JA, van Wijnen M, Stern DM, Nawroth PP, Sixma JJ, Bouma BN. vWF as a regulator of intrinsic factor X activation. *J Lab Clin Med* 1994; 123: 585-93.
67. Neutenschwander PF, Jesty J. Thrombin-activated and factor Xa activated human factor VIII: Differences in cofactor activity and decay rate. *Archs Biochem Biophys* 1992; 296: 426-34.
68. Finke WA, Yu M-Y. W. Characterization of vWF in factor VIII concentrates. *Amer J Hematol* 1989; 31: 41-5.
69. Mazzurri C. Purification, Biochemistry, and Structure/Function Relationships of vWF. In: Factor VIII-vWF. Volume I. Biochemical, Methodological and Functional Aspects. Seghatchian MJ, Savidge GF, eds. Boca Raton, FL: CRC Press 1989; 41-81.
70. Seghatchian MJ. Regulatory Control, Standardization and Characterization of Clinical Concentrates. In: Factor VIII - vWF. Volume II. Clinical Aspects of Deficiency States. Seghatchian MJ, Savidge GF, eds. Boca Raton, FL: CRC Press 1989; 223-53.
71. Aronson DL, Chang P. Ultracentrifugal analysis of factor VIII and von Willebrand Factor in therapeutic preparations. *Vox Sang* 1995; 69: 8-13.
72. Aronson DL, Chang P. Activated factor VIII in therapeutic preparations. Analysis by ultracentrifugation. *Vox Sang* 1996; 71: 142-9.
73. Brandt JT, Triplett DA. Laboratory Assays for Factor VIII. In: Coagulation and Bleeding Disorders. The Role of Factor VIII and vWF. Zimmerman TS, Ruggeri ZM, eds. New York: Marcel Dekker 1989; 343-57.
74. Biggs R, Eveling J, Richards G. The assay of antihemophilic globulin. *Brit J Haematol* 1955; 1: 20-34.
75. Barrowcliffe T. Methodology of the two-stage assay of factor VIII (VIII:C). *Scand J Haematol* 1986; 33 (Suppl. 41): 25-38.
76. Rosen S. Assay of factor VIII with a chromogenic substrate. *Scand J Haematol* 1984; 33 (Suppl. 40): 139-45.
77. van Diejen G, van-Diejen-Visser MP, Franssen J, Hemker HC. Spectrophotometric method for the assay of human blood coagulation factor VIII. *Haemostasis* 1987; 17: 14-24.
78. Hirst CF, Hewitt J, Poller L. Traces of factor VIII in substrate plasma. *Brit J Haematol* 1992; 81: 305-6.
79. Munn KG, Jenny RJ, Krishnaswamy S. Cofactor proteins in the assembly and expression of blood clotting enzyme complexes. *Ann Rev Biochem* 1988; 57: 915-56.
80. Silverberg M. The Assay of Coagulation Factors Using Synthetic Chromogenic Substrates. In: Methods in Hematology, Vol. 7. Disorders of Thrombin Formation. Colbran RW, eds. New York: Churchill Livingstone 1983; 1-23.
81. Andersson M, Andras M, Andersson N-E, Rosen S. Coamate[®] FVIII - a new kit for the accurate chromogenic determination of FVIII activity in plasma and concentrates on automated instruments or manually. *Thromb Haemost* 1993; 9: 1121 (abstract no. 2066).
82. Lang H, Beresdter M, Moritz B. A new chromogenic factor VIII-assay based on a system of human proteins. *Thromb Haemost* 1991; 65: 943 (abstract no. 882).
83. Lee C, Barrowcliffe T, Bray G, Gomperts E, Hubbard A, Kembell-Cook G, Lilly P, Owens D, van Tilburg L, Pasi J. Pharmacokinetics in vivo comparison of using 1-stage and chromogenic substrate assays with two formulations of Hemofil[®] M. *Thromb Haemost* 1996; 76: 950-6.
84. Nilsson IM, Kirkwood TBL, Barrowcliffe TW. In vivo recovery of factor VIII. A comparison of one-stage and two-stage assay methods. *Thromb Haemost* 1979; 42: 1230-9.
85. Hellstern P, Kiehl R, Myashita C, Scherdt H, von Blohn G, Köhler M, Büttner M, Wenzel E. Factor VIII:C (FVIII:C) recovery and half-life after infusion of steam-treated high-purity factor VIII concentrate in severe hemophilia A. A comparison of one-stage assay, two-stage assay and a chromogenic substrate assay. *Thromb Haemost* 1986; 56: 353-9.
86. Messori A, Morfini M, Blomback M, Cinotti S, Longo G, Schimpf K, Schumacher K, Novakova-Banet A, Delvos U, Kjellman H. Pharmacokinetics of two pasteurized factor VIII concentrates by different and multi-center assays of factor VIII activity. *Thromb Res* 1992; 65: 699-708.
87. Blomback M, Kjellman H, Allan JP, Hedner U, Schimpf K, Weichel B. On the unreliability of one-stage factor VIII:C clotting assays after infusion of factor VIII concentrates. *Scand J Clin Lab Invest* 1987; 47: 561-6.
88. Barrowcliffe TW. Standardization and Assay. *Sem Thromb Hemost* 1993; 19: 73-9.
89. Ofosu FA, Blajchman MA, Barr R, Inwood M. Approaches to the Measurement of the In Vivo Response to Factor VIII Concentrate Infusion. *Thromb Res* 1985; 42: 471-6.
90. Mucciari M, Messori A, Donati-Cori G, Longo G, Vannini S, Morfini M, Tendi E, Rossi-Ferrini PL. Kinetic evaluation of four factor VIII concentrates by model-independent methods. *Scand J Haematol* 1985; 34: 22-8.
91. Longo G, Mucciari M, Messori A, Morfini M, Rossi-Ferrini PL. Pharmacokinetics of a new heat-treated concentrate of factor VIII estimated by model-independent methods. *Thromb Res* 1986; 42: 471-6.
92. White GC II, McMillan CW, Kingdon HS, Shoemaker CB. Use of recombinant antihemophilic factor in the treatment of two patients with classic hemophilia. *New Eng J Med* 1989; 320: 166-70.

93. Schwartz RD et al. Human recombinant DNA-derived antihemophilic factor (factor VIII) in the treatment of hemophilia A. *New Eng J Med* 1990; 323: 1800-5.
94. Morfini M, Lee M, Messori A. The design and analysis of half-life and recovery studies for factor VIII and factor IX. *Thromb Haemost* 1991; 66: 384-6.
95. Morfini M, Mannucci PM, Longo G, Cinotti S, Messori A. Comparative evaluation of the pharmacokinetics of three monoclonal factor VIII concentrates. *Thromb Res* 1991; 61: 285-90.
96. Addiego JE, Gomperts E, Liu S-L, Bailey P, Courter SG, Lee ML, Neslund GG, Kingdon HS, Griffith MJ. Treatment of hemophilia A with a highly purified factor VIII concentrate prepared by anti-FVIIIc immunofluorescence chromatography. *Thromb Haemost* 1992; 67: 19-27.
97. Manno CS, Butler RB, Cohen AR. Low recovery in vivo of highly purified factor VIII in patients with hemophilia. *J Pediatr* 1992; 12: 814-8.
98. Björkman S, Carlsson M, Bernlöf E, Stenberg P. Pharmacokinetics of factor VIII in humans. Obtaining clinically relevant data from comparative studies. *Clin Pharmacokinet* 1992; 22: 385-95.
99. Morfini M, et al. Pharmacokinetic properties of recombinant factor VIII compared with a monoclonally purified concentrate (Hemofil M®). *Thromb Haemost* 1992; 68: 433-5.
100. Knvellman A, de Wit HJC, Potstra P, Does JA v.d. Development and small-scale production of a severely heated factor VIII concentrate. *Vox Sang* 1994; 66: 89-95.
101. Woodrow IE, Manson GA. A general expression for coupled enzyme assays of hysteretic enzymes. *Biochem J* 1984; 220: 173-8.
102. Brooks SP, Stuelter CH. Lagtime, a program for calculating coupled enzyme assay parameters. *Intern J Biomed Comput* 1986; 19: 101-7.
103. Yang SY, Schultz H. Kinetics of coupled enzyme reactions. *Biochemistry* 1987; 26: 5579-84.
104. Sergienko EA, Srivastava DK. A continuous spectrophotometric method for the determination of glycogen phosphorylase-catalyzed reaction in the direction of glycogen synthesis. *Analyst Biochem* 1994; 221: 348-55.
105. Barrowcliffe TW, Raut A, Hubbard AR. Discrepancies in potency assessment of recombinant FVIII concentrates. *Haemophilia* 1998; 4: 634-40.
106. Lind P, Larsson K, Spira J, Sydow-Backman M, Almstedt A, Gray E, Sandberg H. Novel forms of B-domain-deleted recombinant factor VIII molecules. Construction and biochemical characterization. *Eur J Biochem* 1995; 232: 19-27.
107. Fatouros A, Osterberg T, Mikaelsson M. Recombinant factor VIII SQ - inactivation kinetics in aqueous solution and the influence of disaccharides and sugar alcohols. *Pharmaceutical Research* 1997; 14: 1679-84.
108. Osterberg T, Fatouros A, Mikaelsson M. Development of freeze-dried albumin-free formulation of recombinant factor VIII SQ. *Pharmaceutical Research* 1997; 14: 892-8.
109. Lusher JM, Hillman-Wiesman C, Horst D. In vivo recovery with products of very high purity - assay discrepancies. *Haemophilia* 1998; 4: 641-5.
110. Mikaelsson M, Oswaldsson U, Sandberg H. Influence of phospholipids on the assessment of factor VIII activity. *Haemophilia* 1998; 4: 646-50.
111. Pittman DP, Alderman EM, Tomkinson KW, Wang JH, Giles AR, Kaufmann RJ. Biochemical, immunological, and in vivo functional characterization of B-domain deleted factor VIII. *Blood* 1993; 81: 2925-35.
112. Merens K, Donath MJS, van Leeuwen RW, de Keyser-Nellen MJM, Verbeet MPH, Klaase Bus JM, Leyte A, van Mourik JA. Biological activity of recombinant factor VIII variants lacking the central B-domain and the heavy-chain sequence Lys713-Arg740: discordant in vitro and in vivo activity. *Br J Haematol* 1993; 85: 133-42.
113. Finney DJ. *Statistical Method in Biological Assay*, 3rd Edition, New York: Macmillan Publishing Co., inc., 1978.

Received June 22, 1999 Accepted after resubmission August 3, 2000